

Ulrich

Foelsche

Institute of Physics/Institute for Geophysics, Astrophysics, and Meteorology (IGAM) and Wegener Center for Climate and Global Change (WEGC), University of Graz, Austria

Lucas Oppeneiger, Institute of Physics/Institute for Geophysics, Astrophysics, and Meteorology (IGAM) and Wegener Center for Climate and Global Change (WEGC), University of Graz, Austria

Julia Danzer, Institute of Physics/Institute for Geophysics, Astrophysics, and Meteorology (IGAM) and Wegener Center for Climate and Global Change (WEGC), University of Graz, Austria

Oral

Atmospheric Rivers (AR) are comparatively narrow regions in the atmosphere that are responsible for most of the horizontal transport of water vapor in the extratropics. ARs are responsible for many extreme precipitation events and floodings at mid-latitudes, including Europe and the US.

While the integrated water vapor content of ARs can be well measured with microwave and infrared sounders, the vertical structure is less well known. We studied if GNSS radio occultation (RO) data are suited to detect atmospheric rivers, using RO data processed with the WEGC occultation processing system, version 5.6.

By analyzing differences between daily values and long-term averages, we can clearly see the temporal evolution and vertical structure of Atmospheric Rivers.

OSTS session

Regional and Global CAL/VAL for Assembling a Climate Data Record

Meeting name

Joint COSMIC Tenth Data Users' Workshop and IROWG-6 Meeting

[Download to PDF](#)