

Gabriel

Cerveira

Federal University of Maranhão

Cláudia Parise, Federal University of Maranhão

Adilson Machado, Federal University of Maranhão

Luciano Pezzi, National Institute for Space Research

Kevin Hodges, University of Reading

Ronald de Souza, National Institute for Space Research

Oral

The increase in anthropogenic emissions to the atmosphere has been implicated in changes of some

climatic phenomena in the coupled Earth system, including the Atlantic Meridional Overturning

Circulation (AMOC). The Tropical Atlantic is a key region for the AMOC, since the North Brazil

Current (NBC) transports the largest amount of heat to the North Atlantic, either through its mean

flow, or through the NBC retroflection eddies. This study aims to assess the changes to the AMOC

and NBC eddies in a CMIP5 future scenario, through climate modelling based on ocean dynamical

downscaling with the Regional Ocean Modelling System (ROMS), using output from two CMIP5

experiments (Historical and RCP8.5) of the Brazilian Earth System Model (BESM). The grid was

set for the 25°S-25°N and 60°W-30°E domain, the experiments were simulated for 20 years each

(Historical: 1986-2005, RCP8.5: 2081-2100), with daily outputs. To identify and track the ocean

eddies, an automated algorithm (TRACK) was applied to Sea Surface Height anomalies. When

compared with time series and vertical profiles of ocean salinity and temperature from PIRATA

buoys, the ROMS Historical experiment simulates these variables very well, especially the

temperature. The AMOC simulated by ROMS presented a very accurate vertical structure, and its

index presented mean transports of 16.9 Sv (Historical) and 13.8 Sv (RCP8.5), suggesting an 18% weakening in the future climate. Wavelet analysis showed that this weakening of the AMOC index peaks at both the six months and the four years periods. The NBC eddies simulated by the ROMS experiments were present predominantly in the austral summer, with very little decrease of eddies frequency in the RCP8.5 scenario, when compared to the changes found for AMOC. These results show that the NBC eddies can play a major role in the thermohaline circulation, by maintaining the transport even under a weakening AMOC tendency.

Presentation file

[cerveira-presentation.pdf](#)

[Download to PDF](#)