

Emilia

Kilpua

University of Helsinki, Finland

"E. Kilpua<sup>1</sup>, S. Pal<sup>1</sup>, S. Good<sup>1</sup>, M. Ala-Lahti<sup>1</sup>, B. Lynch<sup>2</sup>, A. Osmane<sup>1</sup>, E. Palmerio<sup>2</sup>, J. Räsänen<sup>1</sup>, L.-L. Zhao<sup>3</sup>, S. Bale<sup>2</sup>, and M. Stevens<sup>5</sup>

<sup>1</sup>University of Helsinki, Finland

<sup>2</sup>University of California, Berkeley, California, US

<sup>3</sup>University of Alabama, Huntsville, US

<sup>4</sup>Smithsonian Astrophysical Observatory, Cambridge, Massachusetts, US"

Oral

This presentation discusses two interplanetary CMEs (ICMEs) that were observed by PSP when it was at the heliospheric distance  $\sim 0.5$  au from the Sun. The first event occurred on 15-16 March 2019 and the second 25-26 June, 2020. Both ICMEs were related to slow streamer blow-out CMEs, but they interacted in an interesting manner with their surroundings. The March 2019 CME was preceded by a pair of relatively weak shocks and a prominent sheath region. We discuss the formation of magnetic field fluctuations in the sheath and their connection to upstream solar wind structures. The June 2020 CME did not drive a shock, but it nevertheless disturbed the solar wind and heliospheric current sheet/plasma sheet structure ahead, warping it and deflecting the preceding interplanetary field. This interaction also significantly eroded the magnetic flux from the ICME flux rope.

Presentation file

[kilpua-presentation.pdf](#)

YouTube link

<https://youtu.be/jND4Djil7tM?t=311>

[Download to PDF](#)