

Xiangning
Chu

Laboratory for Atmospheric and Space Physics, University of Colorado Boulder, Boulder, Colorado, USA

Jacob Bortnik, Department of Atmospheric and Oceanic Sciences, University of California, Los Angeles, California, USA

Wen Li, Center for Space Physics, Boston University, Boston, Massachusetts, USA

Xiao-Chen Shen, Center for Space Physics, Boston University, Boston, Massachusetts, USA

Qianli Ma, Department of Atmospheric and Oceanic Sciences, University of California, Los Angeles, California, USA

Donglai Ma, Department of Atmospheric and Oceanic Sciences, University of California, Los Angeles, California, USA


David Malaspina, Laboratory for Atmospheric and Space Physics, University of Colorado Boulder, Boulder, Colorado, USA

Sheng Huang, Center for Space Physics, Boston University, Boston, Massachusetts, USA

Poster

Real-world data sets often exhibit imbalanced distributions, which have significantly more data or observations in a specific range of values than the other ranges. For example, space physics data sets, such as geomagnetic indices, relativistic electron fluxes in Earth's radiation belt, and the occurrence and amplitude of solar flares, are typically imbalanced. This is the too-often-too-quiet challenge, one of the fundamental problems in space physics and space weather, and is also a general problem in machine learning. For example, the electron density and plasma fluxes in the Earth's radiation belts can be accurately modeled in our previous studies [Bortnik et al., 2016, 2018; Chu et al., 2017a,b; 2021; Ma et al., 2022a, b]. However, the ML-based models of the plasma waves are usually biased due to the too-often-too-quiet problem both in numerical simulations and observations [Ma et al., 2018; Camporeale et al., 2019; Guo et al., 2021].

We developed a method to solve this problem and applied it to the whistler-mode chorus waves in the Earth's radiation belt. The ML-based wave model used a neural network approach, which takes geomagnetic indices as input and prediction the wave power. As a result, the model can predict not only the quiet time values but also large events. The fact demonstrates that the model provides reliable and stable predictions when the too-often-too-quiet problem is solved. This method of imbalanced regression has wide applications in space physics/weather and a wider field of machine learning techniques.



Poster PDF

[Chu-Xiangning.pdf](#)

Poster category

Geospace/Magnetosphere Research and Applications

Meeting homepage

[Space Weather Workshop 2023](#)

[Download to PDF](#)