


Harry  
Greatorex  
Queen's University Belfast  
Ryan Milligan, Queen's University Belfast  
Phillip Chamberlin, Laboratory for Atmospheric and Space Plasma  
Poster

The chromospheric Lyman-alpha line of neutral hydrogen ( $\text{Ly}\alpha$ ; 1216 Å) is the most intense emission line in the solar spectrum and is believed to constitute a considerable portion of the total radiated energy in solar flares. Here, we present a multi-wavelength study of three M3 flares that were simultaneously observed by RHESSI, GOES, and SDO. Despite having identical X-ray magnitudes these flares show significantly different  $\text{Ly}\alpha$  responses. The peak  $\text{Ly}\alpha$  enhancements above quiescent background for these flares were 1.5 %, 3.3 %, and 6.4 %. However, the predicted  $\text{Ly}\alpha$  enhancements from FISM2 were consistently  $<2$  %. Examining the properties of the nonthermal electrons derived from spectral analysis of HXR observations, an association was found between  $\gamma$  and the peak irradiance enhancements in  $\text{Ly}\alpha$ . Finally, the percentage of nonthermal energy radiated in the  $\text{Ly}\alpha$  line during the impulsive phase was found to range from 1–5 %. Comparatively, the radiative losses in He II (304 Å) were found to range from 0.4–0.9 %. These results may have significant implications for space weather studies and atmospheric modelling, and will influence the interpretation of flare-related  $\text{Ly}\alpha$  observations in Solar Cycle 25.



Poster PDF

[Greatorex-Harry.pdf](#)

Meeting homepage

[Space Weather Workshop 2023](#)

[Download to PDF](#)