


Arfa  
Mubashir  
Georgia State University  
Ashwin Ashok , Georgia State University  
Ernesto Potdevin, Georgia State University  
Enosh Herath Mudiyansele, Georgia State University  
Tharindu Hettiarachchi, Georgia State University  
Viacheslav M Sadykov, Georgia State University  
Xiaochun He, Georgia State University

Poster

Variations in solar activity cause cosmic ray flux changes at different temporal scales and geositions. Our aim is to determine the correlation and cross-correlation time lag between various space parameters and cosmic ray muon counts from two detectors installed at different geomagnetic positions (one in Atlanta, Georgia and the other at Mt Wilson, California, USA). These two detectors are part of the worldwide network of low-cost and portable cosmic ray muon detectors which is under development at Georgia State University for monitoring the space and terrestrial weather at a global scale. To validate our findings, we compare the muon flux measurements with measurements from the Oulu neutron monitor at low cutoff rigidity. The parameters under consideration include sunspot number, solar wind plasma, the Bz component of the interplanetary magnetic field, the Kp index, and the Dst index. Our results indicate significant correlations between cosmic ray intensities at different geomagnetic cutoff rigidities and the space activity parameters. Additionally, our time-lag analysis, employing cross-correlation techniques, aligns well with previously reported findings.



Poster PDF

[Mubashir-Arfa.pdf](#)

Poster category

Geospace/Magnetosphere Research and Applications

Poster session day

Thursday, April 18, 2024

Poster location

16

Meeting homepage

[Space Weather Workshop 2024](#)

[Download to PDF](#)