

Michael
Weimer

Institute of Environmental Physics, University of Bremen, Germany

Michael Hilker, Institute of Environmental Physics, University of Bremen, Germany

Blanca Fuentes Andrade, Institute of Environmental Physics, University of Bremen, Germany

Stefan Noël, Institute of Environmental Physics, University of Bremen, Germany

Max Reuter, Institute of Environmental Physics, University of Bremen, Germany

Michael Buchwitz, Institute of Environmental Physics, University of Bremen, Germany

Heinrich Bovensmann, Institute of Environmental Physics, University of Bremen, Germany

John P. Burrows, Institute of Environmental Physics, University of Bremen, Germany

Hartmut Bösch, Institute of Environmental Physics, University of Bremen, Germany

Oral

The Fast atmospheric traCe gAs retrievaL (FOCAL) algorithm is a radiative transfer code and retrieval scheme for the column-averaged dry-air mole fractions of carbon dioxide (XCO₂) and other trace gases approximating multiple scattering by a single scattering layer whose altitude is part of the retrieved parameters. This makes FOCAL fast and suitable to be applied to large datasets and it has been applied successfully to operating satellite missions such as GOSAT, GOSAT-2 and the Orbiting Carbon Observatory (OCO) version 2. It will also become one of the operational algorithms for the future Anthropogenic CO₂ Monitoring (CO2M) mission. Here, we show the setup of FOCAL when retrieving XCO₂ from level 1b data by OCO-3 Snapshot Area Maps and show first estimates of the CO₂ emissions for specific emission targets in Europe. In the context of preparing for CO2M, this activity is being carried out as part of the German national Integrated Greenhouse Gas Monitoring System (ITMS), for which an operational data assimilation system for greenhouse gases is being set up for Germany and which is one of Germany's responses to the Paris Agreement from 2015.

Meeting homepage

[IWGGMS-20 Workshop](#)

[Download to PDF](#)