


Air Pollutant Emissions in Africa



Eloise Marais with Christine Wiedinmyer, David Pfothenauer, Evan Coffey, Michael Hannigan

AQ in Africa Workshop
8 June 2021





Rapid changes in air pollution in African cities

Trends in NO_2 and NH_3



Combined effect of population growth and air quality degradation



Indicates substantial increase in air pollution exposure in cities in Africa

[Vohra et al., *in prep*]

Crucial that we better characterize sources contributing to air quality degradation across Africa

Emissions are mostly from inefficient combustion

Trash & e-waste burning



Kerosene



Motorbikes



Charcoal production



Cars & congestion



Solid Biofuels



Flaring



Artisanal Oil Refining



Open biomass burning




Generators



Capture many of these with the **DICE-Africa Inventory**

Anthropogenic emissions are diffuse,
but similar in magnitude to emissions from seasonal burning of biomass

DICE-Africa 2006 emissions of NMVOCs and BC




[Marais & Wiedinmyer, 2016]


Solid fuels like charcoal are still dominant energy sources

Charcoal production and use is increasing by **7% per year**, as alternate options like LPG are costly, fluctuate in cost, and supplies are unreliable

Charcoal production, transport and use emissions for 2014



Charcoal production hotspots



Charcoal industry emissions for Africa 2014 may double by 2030

CH₄ emissions, specifically, may outcompete those from open fires in West Africa by 2025



Investment in fossil fuels, despite climate and health impacts

2030 emissions:
all vehicles + power plants
(including powerships)



3-4 likely to be deployed to
South Africa


**Premature deaths due to
future fossil fuel use:**
GEOS-Chem PM_{2.5} and
Vodonos et al. (2018)
concentration-response curve



Total premature deaths in Africa: 48,000

Emissions estimates very sensitive to choice of emission factors

Total and relative emissions from charcoal industry activities




Using more representative OC emission factor from Keita et al. (2018) for charcoal production doubles OC emissions and alters dominant activity

[Bockarie et al., 2020]

Large differences in bottom-up emissions estimates for Africa

Large differences in current emissions estimates leading to wildly different emissions projections



[Marais & Wiedinmyer, 2016]


Size of discrepancies indicative of large uncertainties in anthropogenic activity and emissions factors

Emission Measurements in Africa

Recent work to update emission factors for key sources

Pfotenhauer et al. *ES&T* (2019) updated emission factors for Sub-Saharan Africa




- Trash burning
- Cooking (Commercial)
- Household cooking
- Kerosene lanterns
- Back-up generators



Differences in Ghanaian CO, EC, and OC emissions in kilotonnes between prior and updated emission factors

Evaluation of Emission Inventories

Measurements in Accra and Northern Ghana for source attribution



Sources of EC and OC at roadside sites across Accra, Ghana

Critical Research Needs

- Still large uncertainties regarding general refuse and e-waste burning
- Activity data is typically at the country level, but higher resolution data is needed
- Better guidance is needed regarding temporal variability for input of emissions to models
- Ongoing field and lab measurements are crucial for improved representation of emissions for conditions in Africa
- Need a more extensive network of surface measurements to evaluate and constrain emissions
- Ideally, emissions inventory development should be conducted by individual countries or regions, but this is costly

Other groups developing emissions inventories for Africa

African Anthropogenic Emissions Inventory for gases and particles from 1990 to 2015:

<https://doi.org/10.25326/56> (**Cathy Liousse's group**)

Global CEDS_{GBD-MAPS} inventory uses DICE-Africa as constraints for emissions from 1970 to 2017:

<https://doi.org/10.5281/zenodo.3865670> (**Randall Martin's group**)

Relevant Emissions Inventory Resources

Links to relevant datasets:

Inventory of charcoal industry emissions in Africa for 2014 and 2030:

<https://doi.org/10.5522/04/14595315.v1>

Surface concentrations of PM2.5 in Africa in 2012 and 2030 from GEOS-Chem:

<https://doi.org/10.5522/04/14595729.v1>

DICE-Africa inventory emissions for 2006 and 2013:

<https://doi.org/10.5522/04/14595723.v1>

Satellite-derived isoprene emissions in Africa from OMI and GEOS-Chem:

<https://doi.org/10.5522/04/14595735.v1>

Any Queries? Please contact us:

Eloise Marais: e.marais@ucl.ac.uk

Mike Hannigan: michael.hannigan@Colorado.EDU

Christine Wiedinmyer: Christine.Wiedinmyer@Colorado.EDU